MATH 5A - SAMPLE FINAL EXAM

(1) Find the following limits if they exist. If not, why not?

(a)
$$\lim_{x \to 0^{-}} \frac{|x|}{x} \cos x$$
 (b) $\lim_{x \to 2} \frac{2-x}{x-2}$ (c) $\lim_{x \to \infty} \frac{x-3}{x^2}$

(c)
$$\lim_{x \to \infty} \frac{x-3}{x^2}$$

- Use the difference quotient and definition of derivative to find f'(x) if $f(x) = x^3 x$. (2)
- Find the derivative of each of the following functions and simplify your answer: (3)

(a)
$$f(x) = \sqrt{x} (x^2 + 2)$$

(a)
$$f(x) = \sqrt{x} (x^2 + 2)$$
 (b) $h(x) = (1 + \tan^2 x)^3$ (c) $g(x) = \frac{x}{\sqrt{x^2 + 1}}$

- (4) Find the y-intercept of the line tangent to the curve $y = x \sin x$ at $(\pi, 0)$.

(a)
$$\int_{0}^{\pi/4} \sin x \cos^3 x \ dx$$

(b) $\int_{0}^{2} (3-x)^2 dx$ (c) $\int \frac{x^3}{\sqrt{x^2-1}} dx$

$$\text{(c) } \int \frac{x^3}{\sqrt{x^2 - 1}} \, dx$$

- (6) Given $f(x) = x(1-x)^{2/5}$,
 - (a) find the interval(s) on which the function f is
 - (i) increasing (ii) decreasing (iii) concave up (iv) concave down
 - (b) find all critical points (c) inflection points (d) find all extrema
 - (e) given the above information, sketch a graph of the above function.
- A person in a rowboat 2 miles from the nearest point on a straight shoreline wishes to reach a (7) house 6 miles farther down the shore. If the person can row at a rate of 3 mi/hr and walk at a rate of 5mi/hr. find the least amount of time required to reach the house. (Show all steps you used to determine minimum is absolute)
- Find the absolute min/max of $f(x) = x-2\cos x$ on the interval $[-\pi,\pi]$. (8)
- (9)
- Find the area of the region bounded by the graph of $y = x^3$, the x axis, x = 2 and x = -2. Find the volume of the solid resulting when the region in the first quadrant bounded by the graphs of (10) $y = 4x^2$ and y = 16 is revolved about the x-axis.
 ** <u>SET UP ONLY</u> - TWO WAYS**
 - (a) cylindrical shells

- (b) disks/washers
- A balloon is rising vertically over a point A on the ground at a rate of 15 ft/sec. A point B on the ground (11)is level with A and is 30 ft. from A. When the balloon is 40 ft. above A, at what rate is its distance from B changing?
- Find the equation of the line through (3,4) which cuts from the first quadrant a triangle of minimum area. (12)
- (13) Does the Mean Value Theorem apply to the given function? If so, find "c". If not, why not?

$$f(x) = \sqrt{2x+1}$$
, [0,4]