
MATH-8 TEST Unit 2	
SAMF 100 points	LE NAME:
This test is in two parts. On part one, you may not use a cal necessary. When you complete part one, you turn it in and not go back to it. You will show all work on the test paper, n	culator; on part two, a (non-graphing) calculator is get part two. Once you have turned in part one, you may o scratch paper is allowed.
PART ONE - NO CALC	JLATORS ALLOWED
(1) Find the following Trig Values, exactly (2 point	ts each)
$\cos(7\pi/6) = _ \cot(\pi/3) = _$	$\tan(5\pi/6) =$
$\sin(3\pi/4) = $ $\cos(17\pi/6) = $	$\cos(\pi) =$
$\tan(\pi/2) = _ \sin(-2\pi/3) = _$	$\sin(3\pi/2) = $
$\sin(5\pi/3) = ___\tan(4\pi/3) = _$	$\tan(\pi) = \underline{\qquad}$
(2) Solve the following equations for the given restrict	ion on t. (If no restriction is given, find all solutions) (4 points each)
(a) Solve: $\sin(t) = \frac{\sqrt{2}}{2}$	
(b) Solve: $\cos(t) = -\frac{1}{2}$ for $0 \le t < 2\pi$	
(c) Solve: $\tan(t) = \sqrt{3}$	
(d) Solve: $\sin(t) = 1$	
(e) Solve: $\sec(t) = 2$ for $0 \le t < 2\pi$	
(2) Use the figure to	(1 points each)
(a) approximate the value of sin 5	cos 2
(b) find a value of t such that cost \approx -0.8	$\begin{array}{c} 2.00 \\ 2.25 \\ 0.8 \\ 0.8 \\ 0.75$
(c) find a value of t such that sint $\approx~0.4$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

	NAME:
MATH 8 Sample T	
PART TWO - CALCULATORS ALLOW Show your work on this paper. EXACT answers are expecte graphs and label highs and lows. Give units i	d unless otherwise specified. Show scales on
Fill in the blanks. (2 points each)	
(1) $f(t) = \sin(t)$ Is even, odd, or neither	_
(2) What is the amplitude of $f(t) = -\frac{1}{2}\sin(3t+\pi) - 4$?	
(3) What is the range of $f(t) = \sin(t)$?	
(4) In which quadrant, if any, is $\tan(t) < 0$ and $\sin(t) > 0$ (bo	th true)
(5) The domain of $f(x) = \cot(x)$ is j	
(6) Using your calculator, find approximations for the following, rou (a) $\tan(-3\pi/8) \approx$ (b) $\cos(4) \approx$	
(7) Given $\cos(t) = -\frac{5}{13}$, with <i>t</i> in Quadrant II, find:	(2 points each)
(a) $\sin(t) =$	(b) sec(t)
(8) Given $\tan(t) = -2$ and $\cos(t) < 0$ in, find	(2 points each)

(a) $\sin(t) =$ _____


(b) cos(t)_____


(9) Sketch the following graph. (clearly show scale, graph at least one period, label coordinates of highs and lows) (6 points) $f(x) = -2\sin(3x)$




(10) Sketch the following graph. (clearly show scale, graph at least one period, label coordinates of highs and lows)
(8 points)


 $f(x) = 4\cos\left(\frac{1}{2}x + \frac{\pi}{4}\right)$







(12). Sketch the following graph. (clearly show scale, graph at least TWO periods, If there are any



(13) A mass suspended from a spring is pulled down a distance of 2 feet from its rest position as shown. The mass is released from there at time t=0 and is allowed to oscillate in simple harmonic motion. If the mass returns to this position after 1/3 second, find an equation that describes the motion. (5 points)

